

Immediate Post-Partum LARC Breakout

Maya Oaks, Director of Reproductive Health Programs, ACOG

Nisha Verma, MD Darney-Landy ACOG Fellow

Karena M. Moran, PhD, Research and Quality Project Manager

Northeastern PA Perinatal Quality Collaborative

Name, Title and Organization

PLEASE ADD TO THE CHAT

Addressing Misinformation About Postpartum LARC Through Contraceptive Counseling



ACOG

The American College of
Obstetricians and Gynecologists

the **LARC**
program
Long-Acting Reversible Contraception

ACOG disclaimer

This information is designed as an educational resource to aid clinicians in providing obstetric and gynecologic care, and use of this information is voluntary. This information should not be considered as inclusive of all proper treatments or methods of care or as a statement of the standard of care. It is not intended to substitute for the independent professional judgment of the treating clinician. Variations in practice may be warranted when, in the reasonable judgment of the treating clinician, such course of action is indicated by the condition of the patient, limitations of available resources, or advances in knowledge or technology. The American College of Obstetricians and Gynecologists reviews its publications regularly; however, its publications may not reflect the most recent evidence. Any updates to this document can be found on www.acog.org or by calling the ACOG Resource Center.

While ACOG makes every effort to present accurate and reliable information, this publication is provided "as is" without any warranty of accuracy, reliability, or otherwise, either express or implied. ACOG does not guarantee, warrant, or endorse the products or services of any firm, organization, or person. Neither ACOG nor its officers, directors, members, employees, or agents will be liable for any loss, damage, or claim with respect to any liabilities, including direct, special, indirect, or consequential damages, incurred in connection with this publication or reliance on the information presented.

© 2019 by the American College of Obstetricians and Gynecologists, 409 12th Street SW, PO Box 96920, Washington, DC 20090-6920. Individuals and groups providing patient care or clinical education in family planning have permission to copy all or any portion of this slide set for noncommercial, educational purposes, provided that no modifications are made and proper attribution is given.

Speaker disclosures

- None

Learning objectives

1. Understand key elements of person-centered postpartum contraceptive counseling
2. Discuss common myths and misinformation related to the IUD and implant
3. Apply person-centered postpartum contraceptive counseling to discussions of misinformation with patients

CONTRACEPTIVE COUNSELING

Shared Medical Decision Making

Contraceptive coercion

- **Contraceptive coercion is the act of pressuring or forcing an individual to use a method of birth control that they do not desire**
- **LARC Coercion is the act of forcing or pressuring someone to get or keep a LARC method, or preventing removal of a LARC method**
- **The U.S. has a long history of contraceptive coercion and forced sterilization perpetrated against economically marginalized individuals and persons of color.**

Slide content courtesy of Dr. Serina Floyd

©ACOG



Forced sterilization

- The eugenics movement of the early 1900s
- Continued forced or coerced sterilization through 1970's of the economically marginalized, those with mental illnesses, persons of color, and immigrant individuals
 - Population control
 - Social control
 - Form of punishment
 - Extortion to ensure receipt of public assistance
 - Trainee education
- Recent cases in the 2000s in California prisons, 2020 in Georgia ICE facility


Slide content courtesy of Dr. Serina Floyd

©ACOG

the **LARC**
program
Long-Acting Reversible Contraception

Reproductive injustices

- Mississippi Appendectomy
- Indian Health Services
- La Operación
- Oral contraception clinical trials
- Norplant and Depo Provera



Slide content courtesy of Dr. Serina Floyd

©ACOG

the **LARC**
program
Long-Acting Reversible Contraception

Contraceptive counseling, especially on sterilization or LARC methods, must be sensitive to this history

Slide content courtesy of Dr. Serina Floyd

©ACOG

the **LARC**
program
Long-Acting Reversible Contraception

Reproductive justice

SisterSong defines reproductive justice as:

“The human right to maintain personal bodily autonomy, have children, not have children, and parent the children we have in safe and sustainable communities.”

- **SisterSong Women of Color Reproductive Justice Collective**

A reproductive justice framework for contraceptive counseling

Key Takeaway:

“The framework of reproductive justice connects family planning and other aspects of sexual and reproductive health with the disparities and complexities that affect patients’ lives. Furthermore, it encourages gynecologic health care providers to examine issues of bias and coercion and advocate for equitable access and change.”

- ACOG Committee Opinion #699, Adolescent Pregnancy, Contraception, and Sexual Activity

Provider bias

- **Explicit bias**: a bias that a person is aware of and believes is correct in some manner
- **Implicit bias**: a bias that is unintentional and unconscious but is activated quickly and unknowingly by situational factors
- **Implicit association tests**: <https://implicit.harvard.edu/implicit/>
- **Consequences on patient-provider relationship include:**
 - Rapid discontinuation of methods that client felt pressured to select
 - Delaying future healthcare access and contraceptive use due to previous negative encounters
 - Undermining trust and decreased receptiveness to contraceptive counseling

Slide courtesy of Dr. Serina Floyd

Shared medical decision making for contraceptive counseling

- When engaging in shared medical decision making:
 - Be aware of and address your own biases
 - Practice perspective-taking and individuation when caring for each person
 - Acknowledge historical racial injustices during counseling sessions
 - Strive for equitable outcomes for all people, especially for disadvantaged or marginalized groups.

Talking with patients about contraception

- Shared medical decision making is a process where both patients and clinicians share information, express treatment preferences, and agree on a treatment plan.
- It can increase patient engagement and reduce risk, resulting in improved outcomes, satisfaction, and treatment adherence
- Although medical knowledge is tipped towards the provider, in shared medical decision making a middle ground is sought that incorporates accurate medical information and a patient's personal preferences
- Person-centered goals may also have a part in the decision-making process

©ACOG

Effective patient–physician communication. Committee Opinion No. 587. American College of Obstetricians and Gynecologists. Obstet Gynecol 2014;123:389–93. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24451677>.

Partnering with patients to improve safety. Committee Opinion No. 490. American College of Obstetricians and Gynecologists. Obstet Gynecol 2011;117:1247–9. Available at: http://journals.lww.com/greenjournal/Citation/2011/05000/Committee_Opinion_No_490_Partnering_With.49.aspx.

the **LARC**
program
Long-Acting Reversible Contraception

5 components of shared medical decision making

1.

Focus on interpersonal relationship.

2.

Elicit patient preferences for methods.

3.

Be attuned to diverse patient preferences.

4.

Provide relevant information in accordance with patient preferences.

5.

Be aware of and responsive to patient preferences during counseling.

5 components of shared medical decision making

1. Focus on interpersonal relationship

Intimate, friend-like interactions establish trust and openness between providers and patients and are consistent with patient preferences for counseling about contraception.

- **Examples:**
 - “Hi ____! It’s nice to see you again. How’s everything been since we saw each other last?”
 - “How are you liking the implant you received last time?”

2. Elicit patient preferences for methods

Open the discussion of contraceptive method options with an open-ended question that provides a clear indication that the patients’ preferences are the focus of the discussion.

- **Examples:**
 - “What brings you to our office today?”
 - “What is important to you about your birth control method?”

5 components of shared medical decision making

3. Be attuned to diverse patient preferences

Patients will have varied preferences around issues including the relative importance of preventing pregnancy and the significance of specific side effects, including menstrual changes.

- **Examples:**
 - “It’s really important for me to continue having a monthly period, so it’s less obvious I’m using contraception.”
 - “I absolutely cannot gain any weight.”

4. Provide relevant information in accordance with patient preferences

Prioritize sharing information about methods based on what is most important to the patient, whether that is side effects, efficacy, mode of use, or other method characteristics.

- **Examples:**
 - “Since you said you want regular menses, you might consider oral contraceptive pills.”
 - “I hear you. The injectable contraceptive is the only method proven to cause weight gain, but everybody reacts differently so we can work together to see which method works best for you.”

5 components of shared medical decision making

5. Be aware of and responsive to patient preferences during counseling

Either through direct questioning or by assessing her response to a shared decision making approach, understand and adjust counseling, and specifically the extent of provider guidance in the decision-making process according to how the patient would like decisions to be made.

- **Examples:**
 - “Do you want to use a method that you can easily start and stop on your own?”
 - “How do you feel about having to take a pill at the same time everyday? Does that fit into your daily life?”

Remember the goal! Be person-centered.

Immediate postpartum LARC counseling

- Optimally, patients should be counseled prenatally
- Counseling on immediate postpartum LARC should include:
 - All indicated forms of contraception
 - Advantages, contraindications, and alternatives
 - Increased risk of expulsion, including unrecognized expulsion of IUD
 - Convenience and effectiveness, as well as the benefits of reducing unintended pregnancy and lengthening interpregnancy intervals
 - A discussion on the theoretical risk of reduced duration of breastfeeding
 - Possibility of non-visualized strings and difficult removal
- More info & resources: [ACOG Postpartum Contraceptive Access Initiative Website](#)

ADDRESSING COMMON MYTHS AND MISINFORMATION

Related to LARC Methods

1. Validate concerns/experiences


- I'm sorry your aunt had that experience.
- I'm glad you brought that up.
- Great question. That is a really common concern.
- That sounds frustrating.
- We haven't found that the IUD causes this side effect in studies, but I understand that everyone's body responds differently to medicines, and I'm sorry you are experiencing this.

2. Address misinformation

- **Present accurate information to address myths and misconceptions**
- **Common myths and misconceptions**
 - **The IUD is dangerous – it causes infections and infertility.**
 - **My partner will be able to feel the IUD during sex.**
 - **The implant can move to another part of my body.**
 - **The IUD and implant will make me gain weight.**
 - **The IUD and implant are long-term commitments.**
 - **The IUD and implant will prevent me from being able to breastfeed.**


MYTH: The IUD is dangerous – it causes infections and infertility.

- People did experience problems with previous IUDs that have been off the market for decades (e.g. Dalkon Shield).
- Many studies show no increased risk of pelvic infection or infertility in people using current IUDs.



MYTH: My partner will be able to feel the IUD during sex.

- The IUD is placed at the top of the uterus. Plastic strings are attached to the end and tuck up behind the cervix. They are soft enough that they should not bother your partner.
- If you or your partner are experiencing discomfort during sex, you can go to your healthcare provider to make sure the IUD is in the proper place. The strings can also be adjusted to make you more comfortable if needed.



MYTH: The implant can move to another part of my body.

- The implant can shift a little, but almost always less than an inch from where it started.
- Rarely, if the implant isn't placed correctly, it can end up in a place it shouldn't be.
- You can feel the implant with your fingers if you have any concerns to make sure it is in the right place.



MYTH: The IUD and implant will make me gain weight.

- Overall, studies have not shown an association between the implant and weight gain.
- With the hormonal IUD, there are very low levels of hormones that get into the bloodstream.

MYTH: The IUD and implant are long-term commitments.

- While the IUD and implant prevent pregnancy for many years, you absolutely don't have to keep them in place that whole time.
- You can have either of these methods removed at any time on request.

MYTH: The IUD and implant will prevent me from being able to breastfeed.

- The Copper IUD lacks hormones and is classified as CDC MEC Category 1 (no restriction on use) for people who are breastfeeding
- The LNG IUD and implant are category 2 for theoretical impact on lactation
- Several small randomized control trials (RCTs) have shown no significant differences in:
 - Breast milk quality or quantity
 - Infant size

Legend:			
1	No restriction (method can be used)	3	Theoretical or proven risks usually outweigh the advantages
2	Advantages generally outweigh theoretical or proven risks	4	Unacceptable health risk (method not to be used)

Breastfeeding

Key Takeaway:

“Given available evidence, women considering IPP hormonal LARC should be counseled about the theoretical risks of reduced duration of breastfeeding, but the preponderance of evidence has not shown a negative effect on actual breastfeeding outcomes”

- ACOG Practice Bulletin #186, LARC: IUDs & Implants

3. Be up front about side effects

- **Consider including these tips during counseling so patients can try them before an in-person appointment:**
 - **Copper IUD**
 - **To prevent heavy and painful menses: Take ibuprofen 400mg every 4 hours for 7 days starting Day 1 of menses for the first 3-6 cycles**
 - **LNG-IUD and ENG Implant**
 - **For unscheduled bleeding: Take naproxen 500mg every 12 hours for 5 days OR ibuprofen 800mg every 8 hours for 5 days**

Video: responding to patient concerns

LARC COUNSELING SCENARIOS:

Responding to Patient Concerns

KEY TAKEAWAYS

Things to Keep in Mind

Remember the goal! Be person-centered.

- The goal of contraceptive counseling is to:
 - Discuss all contraceptive options with patients
 - Have patients leave with a plan for contraception they are comfortable with
 - Allow patients to make the contraceptive choices that are best for them
 - Remind patients that there is not one perfect method for everyone
- Shared Medical Decision Making:
 - Focuses on the interpersonal relationship
 - Uses open-ended questions
 - Helps your patient choose the best method for them
 - Takes into account social and cultural contexts

Key takeaways: important things to keep in mind

Respect

- Cultural differences
- Personal preferences
- Dislike of method or side effects
- Patient's decision whether to use birth control after delivery

Listen

- To your patient's preferences
- To your patient's concerns
- Shared decision-making is best, but can take time – be patient

Engage

- Allow patient time to consider their options
- Provide information to help patient make a decision

The ACOG LARC Program can help!

- Email us: pcai@acog.org
- Find more resources online:
 - <https://pcainitiative.acog.org>
 - <https://www.acog.org/programs/long-acting-reversible-contraception-larc>
- Send us your LARC-related questions:
 - www.acoglarc.freshdesk.com
 - The LARC Program Help Desk is a free service open to all, ACOG members and non-members alike
 - All questions will be responded to within 10 business days.

ACOG guidance on contraceptive counseling

ACOG has many contraceptive counseling resources, including, but not limited to:

1. [ACOG Practice Bulletin #186, LARC: Implants and Intrauterine Devices](#)
2. [ACOG Committee Opinion #672, Clinical Challenges of LARC Methods](#)
3. [ACOG Committee Opinion #670, Immediate Postpartum LARC](#)
4. [ACOG Committee Opinion #699, Adolescent Pregnancy, Contraception, and Sexual Activity](#)
5. [ACOG Committee Opinion #490, Partnering With Patients to Improve Safety](#)
6. [ACOG Committee Opinion #587, Effective Patient-Physician Communication](#)
7. [ACOG Committee Opinion #736, Optimizing Postpartum Care](#)
8. [Obstetric Care Consensus #8: Interpregnancy Care](#)

ACOG guidance on care for underserved communities

1. [ACOG Committee Opinion #729, Importance of Social Determinants of Health and Cultural Awareness in the Delivery of Reproductive Health Care](#)
2. [ACOG Committee Opinion #627, Health Care for Unauthorized Immigrants](#)
3. [ACOG Committee Opinion #576, Health Care for Homeless Women](#)
4. [ACOG Committee Opinion #547, Health Care for Women in the Military and Women Veterans](#)
5. [ACOG Committee Opinion #586, Health Disparities in Rural Women](#)
6. [ACOG Committee Opinion #515, Health Care for Urban American Indian and Alaska Native Women](#)
7. [ACOG Committee Opinion #708, Improving Awareness of and Screening for Health Risks Among Sex Workers](#)
8. [ACOG Committee Opinion #518, Intimate Partner Violence](#)
9. [ACOG Committee Opinion #554, Reproductive and Sexual Coercion](#)
10. [ACOG Committee Opinion #830, Reproductive Health Care for Incarcerated Pregnant, Postpartum, and Nonpregnant Individuals](#)
11. [ACOG Committee Opinion #820, Breastfeeding Challenges](#)
12. [ACOG Committee Opinion #496, At-Risk Drinking and Alcohol Dependence: Obstetric and Gynecologic Implication](#)
13. [ACOG Committee Opinion #633, Alcohol Abuse and Other Substance Use Disorders: Ethical Issues in Obstetric and Gynecologic Practice](#)

QUESTIONS?

nverma@acog.org

Contraceptive Counseling: Post-Training Questionnaire



1. Open the Camera App on your smartphone.
2. Hold your device so that the QR code appears in the Camera App's viewfinder.
3. A notification should pop up. Tap the notification to open the link.

- OR -

Access the survey through the following weblink:
<https://www.surveymonkey.com/r/HSFMNPN>

References

- Access to postpartum sterilization. Committee Opinion No. 530. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2012;120:212-5. Available at: http://journals.lww.com/greenjournal/Citation/2012/07000/Committee_Opinion_No_530_Access_to_Postpartum.43.aspx. Retrieved September 26, 2017.
DOI: 10.1097/AOG.0b013e318262e354 [doi].
PMID: 22914423.
- ACQUIRE Project, EngenderHealth. The postpartum intrauterine device: participant handbook. New York (NY): ACQUIRE Project; 2008. Available at: https://www.engenderhealth.org/files/pubs/family-planning/PPIUD_Participant-Handbook.pdf. Retrieved October 3, 2017.
- Adolescent pregnancy, contraception, and sexual activity. Committee Opinion No. 699. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2017;129:e142-9. Available at: http://journals.lww.com/greenjournal/fulltext/2017/05000/Committee_Opinion_No_699_Adolescent_Pregnancy.49.aspx. Retrieved September 21, 2017.
DOI: 10.1097/AOG.0000000000002045 [doi].
PMID: 28426620.
- Affleck PJ, Waisel DB, Cusick JM, Van Decar T. Recall of risks following labor epidural analgesia. *J Clin Anesth* 1998;10:141-4. Available at: [http://www.jcafulltextonline.com/article/S0952-8180\(97\)00258-4/fulltext](http://www.jcafulltextonline.com/article/S0952-8180(97)00258-4/fulltext). Retrieved October 27, 2017.
DOI: S0952818097002584 [pii].
PMID: 9524900.
- Aiken AR, Aiken CE, Trussell J, Guthrie KA. Immediate postpartum provision of highly effective reversible contraception. *BJOG* 2015;122:1050-1. Available at: <http://onlinelibrary.wiley.com/doi/10.1111/1471-0528.13306/abstract>. Retrieved September 13, 2017.
DOI: 10.1111/1471-0528.13306 [doi].
PMID: 25626730.
- American Academy of Pediatrics, American College of Obstetricians and Gynecologists. Guidelines for perinatal care. 8th ed. Elk Grove Village (IL): AAP; Washington, DC: American College of Obstetricians and Gynecologists; 2017.
- American College of Obstetricians and Gynecologists. Contraceptive counseling resource digest. Washington, DC: American College of Obstetricians and Gynecologists; 2017. Available at: <https://www.acog.org/-/media/Departments/LARC/ContraceptiveCounselingReplaceable.pdf>. Retrieved September 21, 2017.
- American College of Obstetricians and Gynecologists. Guidelines for women's health care: a resource manual. 4th ed. Washington, DC: American College of Obstetricians and Gynecologists; 2014.
- Bixby Center for Global Reproductive Health, University of California San Francisco. How well does birth control work? San Francisco (CA): UCSF; 2017. Available at: http://beyondthepill.ucsf.edu/sites/beyondthepill.ucsf.edu/files/Tiers_Chart_ENGLISH.pdf. Retrieved September 22, 2017.
- Bryant AS, Haas JS, McElrath TF, McCormick MC. Predictors of compliance with the postpartum visit among women living in healthy start project areas. *Matern Child Health J* 2006;10:511-6. Available at: <https://link.springer.com/article/10.1007%2Fs10995-006-0128-5>. Retrieved September 13, 2017.
DOI: 10.1007/s10995-006-0128-5 [doi].
PMID: 16807794.
- Bulterys M, Jamieson DJ, O'Sullivan MJ, Cohen MH, Maupin R, Nesheim S, et al. Rapid HIV-1 testing during labor: a multicenter study. *JAMA* 2004;292:219-23. Available at: <https://jamanetwork.com/journals/jama/fullarticle/199058>. Retrieved October 27, 2017.
DOI: 10.1001/jama.292.2.219 [doi].
PMID: 15249571.
- Centers for Disease Control and Prevention. Unintended pregnancy prevention. Atlanta (GA): CDC; 2015. Available at: <https://www.cdc.gov/reproductivehealth/unintendedpregnancy>. Retrieved September 21, 2017.

References

- Clinical challenges of long-acting reversible contraceptive methods. Committee Opinion No. 672. American College of Obstetricians and Gynecologists. Obstet Gynecol 2016;128:e69-77. Available at: http://journals.lww.com/greenjournal/fulltext/2016/09000/Committee_Opinion_No_672_Clinical_Challenges_of.54.aspx. Retrieved September 22, 2017.
DOI: 10.1097/AOG.0000000000001644 [doi].
PMID: 27548557.
- Cohen R, Sheeder J, Arango N, Teal SB, Tocce K. Twelve-month contraceptive continuation and repeat pregnancy among young mothers choosing postdelivery contraceptive implants or postplacental intrauterine devices. Contraception 2016;93:178-83. Available at: [http://www.contraceptionjournal.org/article/S0010-7824\(15\)30030-5/fulltext](http://www.contraceptionjournal.org/article/S0010-7824(15)30030-5/fulltext). Retrieved September 13, 2017.
DOI: 10.1016/j.contraception.2015.10.001 [doi].
PMID: 26475368.
- Connolly A, Thorp J, Pahel L. Effects of pregnancy and childbirth on postpartum sexual function: a longitudinal prospective study. Int Urogynecol J Pelvic Floor Dysfunct 2005;16:263-7. Available at: <https://link.springer.com/article/10.1007/s00192-005-1293-6>. Retrieved September 13, 2017.
DOI: 10.1007/s00192-005-1293-6 [doi].
PMID: 15838587.
- Crockett AH, Pickell LB, Heberlein EC, Billings DL, Mills B. Six- and twelve-month documented removal rates among women electing postpartum inpatient compared to delayed or interval contraceptive implant insertions after Medicaid payment reform. Contraception 2017;95:71-6. Available at: [http://www.contraceptionjournal.org/article/S0010-7824\(16\)30145-7/fulltext](http://www.contraceptionjournal.org/article/S0010-7824(16)30145-7/fulltext). Retrieved September 13, 2017.
DOI: S0010-7824(16)30145-7 [pii].
PMID: 27400823.
- Curtis KM, Jatlaoui TC, Tepper NK, Zapata LB, Horton LG, Jamieson DJ, et al. U.S. selected practice recommendations for contraceptive use, 2016. MMWR Recomm Rep 2016;65(RR-4):1-66. Available at: <https://www.cdc.gov/mmwr/volumes/65/rr/rr6504a1.htm>. Retrieved November 2, 2017.
DOI: 10.15585/mmwr.rr6504a1 [doi].
PMID: 27467319.
- Curtis KM, Tepper NK, Jatlaoui TC, Berry-Bibee E, Horton LG, Zapata LB, et al. U.S. medical eligibility criteria for contraceptive use, 2016. MMWR Recomm Rep 2016;65(RR-3):1-103. Available at: https://www.cdc.gov/mmwr/volumes/65/rr/rr6503a1.htm?s_cid=rr6503a1_w. Retrieved September 13, 2017.
DOI: 10.15585/mmwr.rr6503a1 [doi].
PMID: 27467196.
- Dahlke JD, Terpstra ER, Ramseyer AM, Busch JM, Rieg T, Magann EF. Postpartum insertion of levonorgestrel--intrauterine system at three time periods: a prospective randomized pilot study. Contraception 2011;84:244-8. Available at: [http://www.contraceptionjournal.org/article/S0010-7824\(11\)00008-4/fulltext](http://www.contraceptionjournal.org/article/S0010-7824(11)00008-4/fulltext). Retrieved September 13, 2017.
DOI: 10.1016/j.contraception.2011.01.007 [doi].
PMID: 21843688.
- Dehlendorf C, Fox E, Sobel L, Borrero S. Patient-centered contraceptive counseling: evidence to inform practice. Curr Obstet Gynecol Rep 2016;5:55-63. Available at: <https://link.springer.com/article/10.1007/s13669-016-0139-1>. Retrieved October 27, 2017.
DOI: 10.1007/s13669-016-0139-1.

References

- Dehlendorf C, Krajewski C, Borrero S. Contraceptive counseling: best practices to ensure quality communication and enable effective contraceptive use. Clin Obstet Gynecol 2014;57:659-73. Available at: http://journals.lww.com/clinicalobgyn/Abstract/2014/12000/Contraceptive_Counseling_Best_Practices_to.5.aspx. Retrieved November 1, 2017.
DOI: 10.1097/GRF.0000000000000059 [doi].
PMID: 25264697.
- Dehlendorf C, Levy K, Kelley A, Grumbach K, Steinauer J. Women's preferences for contraceptive counseling and decision making. Contraception 2013;88:250-6. Available at: [http://www.contraceptionjournal.org/article/S0010-7824\(12\)00901-8/fulltext](http://www.contraceptionjournal.org/article/S0010-7824(12)00901-8/fulltext). Retrieved September 21, 2017.
DOI: 10.1016/j.contraception.2012.10.012 [doi].
PMID: 23177265.
- Effective patient–physician communication. Committee Opinion No. 587. American College of Obstetricians and Gynecologists. Obstet Gynecol 2014;123:389-93. Available at: http://journals.lww.com/greenjournal/fulltext/2014/02000/Committee_Opinion_No_587_Effective.36.aspx. Retrieved October 3, 2017.
DOI: 10.1097/01.AOG.0000443279.14017.12 [doi].
PMID: 24451677.
- Elwyn G, Frosch D, Thomson R, Joseph-Williams N, Lloyd A, Kinnersley P, et al. Shared decision making: a model for clinical practice. J Gen Intern Med 2012;27:1361-7. Available at: <https://link.springer.com/article/10.1007%2Fs11606-012-2077-6>. Retrieved October 27, 2017.
DOI: 10.1007/s11606-012-2077-6 [doi].
PMID: 22618581.
- Finer LB, Zolna MR. Declines in unintended pregnancy in the United States, 2008-2011. N Engl J Med 2016;374:843-52. Available at: <http://www.nejm.org/doi/full/10.1056/NEJMsa1506575>. Retrieved September 21, 2017.
DOI: 10.1056/NEJMsa1506575 [doi].
PMID: 26962904.
- Godfrey EM, Folger SG, Jeng G, Jamieson DJ, Curtis KM. Treatment of bleeding irregularities in women with copper-containing IUDs: A systematic review. Contraception. 2013;87(5):549-566.
DOI:10.1016/j.contraception.2012.09.006
- Gray RH, Campbell OM, Zaccaro HA, Labbok MH, MacRae SL. Postpartum return of ovarian activity in nonbreastfeeding women monitored by urinary assays. J Clin Endocrinol Metab 1987;64:645-50. Available at: <https://academic.oup.com/jcem/article-abstract/64/4/645/2653643/Postpartum-Return-of-Ovarian-Activity-in>. Retrieved September 26, 2017.
DOI: 10.1210/jcem-64-4-645 [doi].
PMID: 3818896.
- Guttmacher Institute. Unintended pregnancy in the United States. New York (NY): Guttmacher Institute; 2016. Available at: <https://www.guttmacher.org/fact-sheet/unintended-pregnancy-united-states>. Retrieved September 21, 2017.
- Hatcher RA, Trussell J, Nelson AL, Cates WJ, Kowal D, Policar MS. Contraceptive technology. 20th rev. ed. New York (NY): Ardent Media; 2011.
- The Health Foundation. Shared decision making. Available at: <https://www.youtube.com/watch?v=fhiwftNLtyc>. Retrieved October 27, 2017.
- Immediate postpartum long-acting reversible contraception. Committee Opinion No. 670. American College of Obstetricians and Gynecologists. Obstet Gynecol 2016;128:e32-7. Available at: <http://journals.lww.com/greenjournal/pages/results.aspx?txtkeywords=10.1097%2fAOG.0000000000001587>. Retrieved September 13, 2017.
DOI: 10.1097/AOG.0000000000001587 [doi].
PMID: 27454734.

References

- Informed Medical Decisions Foundation. Shared decision making animated short. Available at: <https://www.youtube.com/watch?v=XPm5iEDEl8Y>. Retrieved October 27, 2017.
- Jackson A, Henry R, Avery N, VanDenKerkhof E, Milne B. Informed consent for labour epidurals: what labouring women want to know. *Can J Anaesth* 2000;47:1068-73. Available at: <https://link.springer.com/article/10.1007%2FBF03027957>. Retrieved October 27, 2017.
DOI: 10.1007/BF03027957 [doi].
PMID: 11097535.
- Kapp N, Curtis K, Nanda K. Progestogen-only contraceptive use among breastfeeding women: a systematic review. *Contraception* 2010;82:17-37. Available at: [http://www.contraceptionjournal.org/article/S0010-7824\(10\)00046-6/fulltext](http://www.contraceptionjournal.org/article/S0010-7824(10)00046-6/fulltext). Retrieved September 13, 2017.
DOI: 10.1016/j.contraception.2010.02.002 [doi].
PMID: 20682140.
- Kongnyuy EJ, Mbu ER, Mbopi-Keou FX, Fomulu N, Nana PN, Tebeu PM, et al. Acceptability of intrapartum HIV counselling and testing in Cameroon. *BMC Pregnancy Childbirth* 2009;9:9. Available at: <https://bmcpregnancychildbirth.biomedcentral.com/articles/10.1186/1471-2393-9-9>. Retrieved October 27, 2017.
DOI: 10.1186/1471-2393-9-9 [doi].
PMID: 19250517.
- Kyleena (levonorgestrel-releasing intrauterine system). Highlights of prescribing information. Whippany (NJ): Bayer HealthCare Pharmaceuticals Inc.; 2016. Available at: http://labeling.bayerhealthcare.com/html/products/pi/Kyleena_PI.pdf. Retrieved August 23, 2017.
- Liletta (levonorgestrel-releasing intrauterine system). Highlights of prescribing information. Irvine (CA): Allergan USA, Inc.; 2017. Available at: https://www.allergan.com/assets/pdf/lilettashi_pi. Retrieved August 23, 2017.
- Long-acting reversible contraception: implants and intrauterine devices. Practice Bulletin No. 186. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2017;130:e251-69. Available at: http://journals.lww.com/greenjournal/Fulltext/2017/11000/Practice_Bulletin_No_186_Long_Acting_Reversible.50.aspx. Retrieved October 25, 2017.
DOI: 10.1097/AOG.0000000000002400 [doi].
PMID: 29064972.
- Lopez LM, Bernholz A, Hubacher D, Stuart G, Van Vliet HA. Immediate postpartum insertion of intrauterine device for contraception. *Cochrane Database of Systematic Reviews* 2015, Issue 6. Art. No.: CD003036. DOI: 10.1002/14651858.CD003036.pub3. Available at: <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD003036.pub3/abstract>. Retrieved October 3, 2017.
DOI: 10.1002/14651858.CD003036.pub3.
- McNicholas C, Swor E, Wan L, Peipert JF. Prolonged use of the etonogestrel implant and levonorgestrel intrauterine device: 2 years beyond Food and Drug Administration-approved duration. *Am J Obstet Gynecol* 2017;216:586.e1-6. Available at: [http://www.ajog.org/article/S0002-9378\(17\)30171-0/fulltext](http://www.ajog.org/article/S0002-9378(17)30171-0/fulltext). Retrieved September 21, 2017.
DOI: S0002-9378(17)30171-0 [pii].
PMID: 28147241.
- Mirena (levonorgestrel-releasing intrauterine system). Highlights of prescribing information. Whippany (NJ): Bayer HealthCare Pharmaceuticals Inc.; 2017. Available at: https://labeling.bayerhealthcare.com/html/products/pi/Mirena_PI.pdf. Retrieved August 23, 2017.
- New York City Department of Health and Mental Hygiene. Breastfeeding and birth control. New York (NY): DOHMH; 2016. Available at: <https://www1.nyc.gov/assets/doh/downloads/pdf/ms/breastfeeding-birth-control-postpartum-poster.pdf>. Retrieved September 21, 2017.
- Nexplanon (etonogestrel implant). Highlights of prescribing information. Whitehouse Station (NJ): Merck & Co., Inc.; 2017. Available at: http://www.merck.com/product/usa/pi_circulars/n/nexplanon/nexplanon_pi.pdf. Retrieved August 23, 2017.

References

Optimizing postpartum care. Committee Opinion No. 736. American College of Obstetricians and Gynecologists. Obstet Gynecol 2018;131:e140-50. Available at: https://journals.lww.com/greenjournal/Fulltext/2018/05000/ACOG_Committee_Opinion_No_736_Optimizing.42.aspx. Retrieved June 12, 2018.

DOI: 10.1097/AOG.0000000000002633 [doi].

PMID: 29683911.

ParaGard T 380A intrauterine copper contraceptive. Prescribing information. North Wales (PA): Teva Women's Health, Inc.; 2014. Available at: <http://paragard.com/pdf/PARAGARD-PI.pdf>. Retrieved August 23, 2017.

Partnering with patients to improve safety. Committee Opinion No. 490. American College of Obstetricians and Gynecologists. Obstet Gynecol 2011;117:1247-9. Available at: http://journals.lww.com/greenjournal/Citation/2011/05000/Committee_Opinion_No_490_Partnering_With.49.aspx. Retrieved September 21, 2017.

DOI: 10.1097/AOG.0b013e31821d7d60 [doi].

PMID: 21508784.

Peipert JF, Zhao Q, Allsworth JE, Petrosky E, Madden T, Eisenberg D, et al. Continuation and satisfaction of reversible contraception. Obstet Gynecol 2011;117:1105-13. Available at: http://journals.lww.com/greenjournal/fulltext/2011/05000/Continuation_and_Satisfaction_of_Reversible.12.aspx. Retrieved September 21, 2017.

DOI: 10.1097/AOG.0b013e31821188ad [doi].

PMID: 21508749.

Postpartum Birth Control. FAQ194. American College of Obstetricians and Gynecologists. 2016. Available at: <https://www.acog.org/Patients/FAQs/Postpartum-Birth-Control>. Retrieved October 25, 2017.

Reilly PR. Eugenics and involuntary sterilization: 1907–2015. Ann Rev Genom Hum Genet 2015;16:351-68. Available at: <http://www.annualreviews.org/doi/abs/10.1146/annurev-genom-090314-024930>. Retrieved September 21, 2017.

DOI: <https://doi.org/10.1146/annurev-genom-090314-024930>.

Rivera R, Yacobson I, Grimes D. The mechanism of action of hormonal contraceptives and intrauterine contraceptive devices. Am J Obstet Gynecol 1999;181:1263-9. Available at: [http://www.ajog.org/article/S0002-9378\(99\)70120-1/fulltext](http://www.ajog.org/article/S0002-9378(99)70120-1/fulltext). Retrieved September 13, 2017.

DOI: S0002-9378(99)70120-1 [pii].

PMID: 10561657.

Rodriguez MI, Kaunitz AM. An evidence-based approach to postpartum use of depot medroxyprogesterone acetate in breastfeeding women. Contraception 2009;80:4-6. Available at: [http://www.contraceptionjournal.org/article/S0010-7824\(09\)00048-1/fulltext](http://www.contraceptionjournal.org/article/S0010-7824(09)00048-1/fulltext). Retrieved September 13, 2017.

DOI: 10.1016/j.contraception.2008.12.014 [doi].

PMID: 19501209.

Sexual assault. ACOG Committee Opinion No. 777. American College of Obstetricians and Gynecologists. Obstet Gynecol 2019;133:e296–302. Available at: <https://www.acog.org/clinical/clinical-guidance/committee-opinion/articles/2019/04/sexual-assault>. Retrieved March 30, 2020.

Shaamash AH, Sayed GH, Hussien MM, Shaaban MM. A comparative study of the levonorgestrel-releasing intrauterine system Mirena versus the Copper T380A intrauterine device during lactation: breast-feeding performance, infant growth and infant development. Contraception 2005;72:346-51. Available at: [http://www.contraceptionjournal.org/article/S0010-7824\(05\)00109-5/fulltext](http://www.contraceptionjournal.org/article/S0010-7824(05)00109-5/fulltext). Retrieved September 13, 2017.

DOI: S0010-7824(05)00109-5 [pii].

PMID: 16246660.

Skyla (levonorgestrel-releasing intrauterine system). Highlights of Prescribing Information. Whippany (NJ): Bayer HealthCare Pharmaceuticals Inc.; 2017. Available at: http://labeling.bayerhealthcare.com/html/products/pi/Skyla_PI.pdf. Retrieved August 23, 2017.

References

- Sothornwit J, Werawatakul Y, Kaewrudee S, Lumbiganon P, Laopaiboon M. Immediate versus delayed postpartum insertion of contraceptive implant for contraception. Cochrane Database of Systematic Reviews 2017, Issue 4. Art. No.: CD011913. DOI: 10.1002/14651858.CD011913.pub2. Available at: <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD011913.pub2/abstract>. Retrieved October 2, 2017.
DOI: 10.1002/14651858.CD011913.pub2.
- Sterilization for Women and Men. FAQ011. American College of Obstetricians and Gynecologists. 2015. Available at: <https://www.acog.org/Patients/FAQs/Sterilization-for-Women-and-Men>. Retrieved December 5, 2017.
- Stern AM. Sterilized in the name of public health: race, immigration, and reproductive control in modern California. Am J Public Health 2005;95:1128-38. Available at: <http://ajph.aphapublications.org/doi/abs/10.2105/AJPH.2004.041608>. Retrieved September 21, 2017.
DOI: 95/7/1128 [pii].
PMID: 15983269.
- Thiel de Bocanegra H, Chang R, Menz M, Howell M, Darney P. Postpartum contraception in publicly-funded programs and interpregnancy intervals. Obstet Gynecol 2013;122:296-303. Available at: http://journals.lww.com/greenjournal/fulltext/2013/08000/Postpartum_Contraception_in_Publicly_Funded.17.aspx. Retrieved September 13, 2017.
DOI: 10.1097/AOG.0b013e3182991db6 [doi].
PMID: 23969798.
- Thurman AR, Janecek T. One-year follow-up of women with unfulfilled postpartum sterilization requests. Obstet Gynecol 2010;116:1071-7. Available at: http://journals.lww.com/greenjournal/fulltext/2010/11000/One_Year_Follow_up_of_Women_With_Unfulfilled.12.aspx. Retrieved September 21, 2017.
DOI: 10.1097/AOG.0b013e3181f73eaa [doi].
PMID: 20966691.
- Trussell J. Contraceptive failure in the United States. Contraception 2011;83:397-404. Available at: [http://www.contraceptionjournal.org/article/S0010-7824\(11\)00049-7/fulltext](http://www.contraceptionjournal.org/article/S0010-7824(11)00049-7/fulltext). Retrieved September 13, 2017.
DOI: 10.1016/j.contraception.2011.01.021 [doi].
PMID: 21477680.
- Voedisch AJ, Blumenthal PD. Postpartum contraception: ways to avoid VTE. Contemp Ob/Gyn 2012;57(1):20-6, 28-9. Available at: <http://contemporaryobgyn.modernmedicine.com/contemporary-obgyn/news/modernmedicine/modern-medicine-feature-articles/postpartum-contraception-way>. Retrieved September 21, 2017.
- Whitaker AK, Chen BA, Borgatta L. Society of Family Planning guidelines: postplacental insertion of intrauterine devices. Contraception 2017; DOI: 10.1016/j.contraception.2017.09.014. Available at: [http://www.contraceptionjournal.org/article/S0010-7824\(17\)30473-0/fulltext](http://www.contraceptionjournal.org/article/S0010-7824(17)30473-0/fulltext). Retrieved October 16, 2017.
DOI: 10.1016/j.contraception.2017.09.014
PMID: 28987293.
- White K, Teal SB, Potter JE. Contraception after delivery and short interpregnancy intervals among women in the United States. Obstet Gynecol 2015;125:1471-7. Available at: http://journals.lww.com/greenjournal/fulltext/2015/06000/Contraception_After_Delivery_and_Short.28.aspx. Retrieved September 21, 2017.
DOI: 10.1097/AOG.0000000000000841 [doi].
PMID: 26000519.
- Woo I, Seifert S, Hendricks D, Jamshidi RM, Burke AE, Fox MC. Six-month and 1-year continuation rates following postpartum insertion of implants and intrauterine devices. Contraception 2015;92:532-5. Available at: [http://www.contraceptionjournal.org/article/S0010-7824\(15\)00582-X/fulltext](http://www.contraceptionjournal.org/article/S0010-7824(15)00582-X/fulltext). Retrieved September 13, 2017.
DOI: 10.1016/j.contraception.2015.09.007 [doi].
PMID: 26408376.

References

World Health Organization. Medical eligibility criteria for contraceptive use. 5th ed. Geneva: WHO; 2015. Available at:

http://www.who.int/reproductivehealth/publications/family_planning/MEC-5/en/. Retrieved September 27, 2017.

Zapata LB, Murtaza S, Whiteman MK, Jamieson DJ, Robbins CL, Marchbanks PA, et al. Contraceptive counseling and postpartum contraceptive use. Am J Obstet Gynecol 2015;212:171.e1-

8. Available at: [http://www.ajog.org/article/S0002-9378\(14\)00808-4/fulltext](http://www.ajog.org/article/S0002-9378(14)00808-4/fulltext). Retrieved October 3, 2017.

DOI: 10.1016/j.ajog.2014.07.059 [doi].

PMID: 25093946.

Zigler RE, McNicholas C. Unscheduled vaginal bleeding with progestin-only contraceptive use. Am J Obstet Gynecol. 2017;216(5):443-450. doi:10.1016/j.ajog.2016.12.008